

<p><b>Active ingredients (generic names)</b></p>	<p><b>IVERMECTIN</b> (i-ver-mec-tin)</p> 
<p><b>What it is used for</b></p>	<ul style="list-style-type: none"> <li>• Crusted scabies</li> <li>• Scabies</li> <li>• Strongyloides</li> </ul>
<p><b>How it works</b></p>	<ul style="list-style-type: none"> <li>• Kills worms and mites that infect people</li> </ul>
<p><b>Side effects</b></p>	<div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;">  <p>Diarrhoea</p> </div> <div style="text-align: center;">  <p>Nausea</p> </div> <div style="text-align: center;">  <p>Sleepy</p> </div> </div> <div style="display: flex; justify-content: space-around; align-items: flex-start; margin-top: 20px;"> <div style="text-align: center;">  <p>Itch</p> </div> <div style="text-align: center;">  <p>Dizzy</p> </div> </div>

 <p><b>Warnings</b></p>	 <p>Pregnant</p>  <p>Children under 5 years or less than 15kg</p>
<p><b>Tell the patient</b></p> 	<ul style="list-style-type: none"> <li>• May need to treat again after 1–2 weeks</li> <li>• How to prevent scabies and/or strongyloides infections</li> <li>• <b>For crusted scabies</b> <ul style="list-style-type: none"> <li>▸ Use skin cream like <i>Lyclear</i> at same time</li> <li>▸ Treat other family members for scabies to prevent reinfection</li> </ul> </li> </ul>
<p><b>Check</b></p>	<p>For strongyloidiasis — do faeces test to see if treatment worked</p>