

Benzathine benzylpenicillin (Bicillin L-A) Inj: 1,200,000units/2.3mL syringe 600,000units/1.2mL syringe Pregnancy: A – safe to use Breastfeed: safe to use	Chickenpox Skin sores Sore throat	Deep IM Single dose	450,000units/0.9mL (337.5mg)	600,000units/1.2mL (450mg) For child 10-19kg	1,200,000units/2.3mL (1 x 2.3mL syringe) (900mg)	Long lasting low levels of penicillin. Do not give for pneumonia. Note: 1,200,000units=900mg For syphilis dose see STI protocols.
	RHD	Deep IM Every 21-28days	600,000units/1.2mL (1 x 1.2mL syringe) (450mg) For child less than 10kg paediatrician consult for secondary prophylaxis regimen		1,200,000units/2.3mL (1 x 2.3mL syringe) (900mg)	

Benzathine benzylpenicillin (Bicillin L-A) Inj: 1,200,000units/2.3mL syringe 600,000units/1.2mL syringe Pregnancy: A – safe to use Breastfeed: safe to use	Chickenpox Skin sores Sore throat	Deep IM Single dose	450,000units/0.9mL (337.5mg)	600,000units/1.2mL (450mg) For child 10-19kg	1,200,000units/2.3mL (1 x 2.3mL syringe) (900mg)	Long lasting low levels of penicillin. Do not give for pneumonia. Note: 1,200,000units=900mg For syphilis dose see STI protocols.
	RHD	Deep IM Every 21-28days	600,000units/1.2mL (1 x 1.2mL syringe) (450mg) For child less than 10kg paediatrician consult for secondary prophylaxis regimen		1,200,000units/2.3mL (1 x 2.3mL syringe) (900mg)	

Benzathine benzylpenicillin (Bicillin L-A) Inj: 1,200,000units/2.3mL syringe 600,000units/1.2mL syringe Pregnancy: A – safe to use Breastfeed: safe to use	Chickenpox Skin sores Sore throat	Deep IM Single dose	450,000units/0.9mL (337.5mg)	600,000units/1.2mL (450mg) For child 10-19kg	1,200,000units/2.3mL (1 x 2.3mL syringe) (900mg)	Long lasting low levels of penicillin. Do not give for pneumonia. Note: 1,200,000units=900mg For syphilis dose see STI protocols.
	RHD	Deep IM Every 21-28days	600,000units/1.2mL (1 x 1.2mL syringe) (450mg) For child less than 10kg paediatrician consult for secondary prophylaxis regimen		1,200,000units/2.3mL (1 x 2.3mL syringe) (900mg)	