

Table 4.5 HIGH cardiovascular risk (if one or more present)

• Known cardiovascular disease (angina, heart attack, bypass surgery, stroke)
• Diabetes <i>AND</i> kidney disease with urine ACR 2.5mg/mmol or more for males, 3.5mg/mmol or more for females
• Diabetes <i>AND</i> age over 60 years
• Chronic kidney disease with eGFR less than 45 or urine ACR more than 25 in males or more than 35 in females
• Persistent high BP — systolic 180mmHg or more <i>OR</i> diastolic 110mmHg or more
• Total cholesterol more than 7.5mmol/L
• Familial hypercholesterolaemia (genetic disorder with high cholesterol)
• 75 years or over

Table 4.5 HIGH cardiovascular risk (if one or more present)

• Known cardiovascular disease (angina, heart attack, bypass surgery, stroke)
• Diabetes <i>AND</i> kidney disease with urine ACR 2.5mg/mmol or more for males, 3.5mg/mmol or more for females
• Diabetes <i>AND</i> age over 60 years
• Chronic kidney disease with eGFR less than 45 or urine ACR more than 25 in males or more than 35 in females
• Persistent high BP — systolic 180mmHg or more <i>OR</i> diastolic 110mmHg or more
• Total cholesterol more than 7.5mmol/L
• Familial hypercholesterolaemia (genetic disorder with high cholesterol)
• 75 years or over

Table 4.5 HIGH cardiovascular risk (if one or more present)

• Known cardiovascular disease (angina, heart attack, bypass surgery, stroke)
• Diabetes <i>AND</i> kidney disease with urine ACR 2.5mg/mmol or more for males, 3.5mg/mmol or more for females
• Diabetes <i>AND</i> age over 60 years
• Chronic kidney disease with eGFR less than 45 or urine ACR more than 25 in males or more than 35 in females
• Persistent high BP — systolic 180mmHg or more <i>OR</i> diastolic 110mmHg or more
• Total cholesterol more than 7.5mmol/L
• Familial hypercholesterolaemia (genetic disorder with high cholesterol)
• 75 years or over